

Beat Drug Fund

ID	Target	Pretest	Intervention	Posttest	Positive outcome
Set No. 1 (Attitude)	Participants	13-item, 5-point Gender, Age		13-item, 5-point	incline to agree after intervention
Set No. 2 (Knowledge)	Participants	10-item, 5-point Gender, Age		10-item, 5-point	Incline to agree after intervention
Set No. 3 (Self-efficacy to avoid drug use)	Participants	16-item, 7-point Gender, Age		16-item, 7-point	Enhance self-efficacy after intervention
Set No. 4 (Self-efficacy to refuse drug use) (English version)	Participants	2 questions 14-item, 5-point Gender, Age		2 questions 14-item, 5-point	Incline to “easy to refuse)
Set No. 5A (Frequency of drug use in the past 30 days) (English version)	Participants	1 question 13-item, 3-point Gender, Age		1 question 13-item, 3-point	Reduced frequency
Set No. 6A (Frequency of drug use in the past 30 months) (English version)	Participants	1 question 13-item, 3-point Gender, Age		1 question 13-item, 3-point	Reduced frequency
Set No. 7A (Frequency of drug use in the past 6 months) (English version)	Participants	1 question 13-item, 3-point Gender, Age		1 question 13-item, 3-point	Reduced frequency
Set No. 8 (Rating for preventive or publicity events)	Participants			5 questions Q1 4-point	Reduced frequency

				Q2 5-point Q3,4 10 item Multi Q5 4-item 5-point Gender, Age	
Set No. 9 (Treatment needs and motivation) (English version)	Participants	36-item, 5-point Gender, Age		36-item, 5-point	Decreased needs Maintained or enhanced motivation
Set No. 10 (Parent's rating on preventive talk or activity)	Parents			4 questions Q1 10-item 4-point Q5 5-point Q6 4-point Q7 9 Gender, Age	Incline to agree on the effectiveness of preventive talk or activity
Set No. 12 (Relapse coping questionnaire) (English version)	Participants	6 questions Q1 4 Fre Q2-6 10-point Gender, Age		6 questions Q1 4 Fre Q2-6 10-point	Enhanced coping skills